

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 859 051 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:

16.01.2002 Bulletin 2002/03

(51) Int Cl.7: **C12N 11/08**

(21) Application number: **98301028.1**

(22) Date of filing: **12.02.1998**

(54) **Immobilized biocatalyst**

Immobilisierter Biokatalysator

Biocatalyseur immobilisé

(84) Designated Contracting States:
DE FR GB

(30) Priority: **14.02.1997 JP 3064797**

(43) Date of publication of application:
19.08.1998 Bulletin 1998/34

(73) Proprietor: **NIPPON SHOKUBAI CO., LTD.**
Chuo-ku, Osaka-shi, Osaka (JP)

(72) Inventors:
• **Mukouyama, Masaharu**
Ibaraki (JP)

• **Komatsuzaki, Satomi**
Ibaraki (JP)
• **Sakano, Koichi**
60439 Frankfurt am Main (DE)

(74) Representative: **Hale, Stephen Geoffrey et al**
JY & GW Johnson, Kingsbourne House, 229-231
High Holborn
London WC1V 7DP (GB)

(56) References cited:
EP-A- 0 446 948 **WO-A-97/46590**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 859 051 B1

Description

BACKGROUND OF INVENTION

1. Field of Invention

[0001] The present invention relates to a novel immobilized biocatalyst and its use.

2. Related Art

[0002] Known immobilized biocatalysts include those wherein enzymes or microorganisms are bound to the surface of a solid support by physical adhesion, ion-bonding or covalent bonding, those wherein enzyme-containing substances, such as cells, are mutually crosslinked, and those wherein enzymes or enzyme-containing substances are covered in a lattice or microcapsules. A different type of immobilized catalyst is described in Japanese Unexamined Patent Publication No. 5-344898, wherein a polyazetidine prepolymer, carboxymethyl cellulose, polyurethane hydrogel prepolymer or polymethylene isocyanate is hardened on the surface of a solid support to immobilize microbial cells on the surface of the support. A disadvantage of these methods, however, is that the enzymes undergo considerable inactivation during the immobilization.

[0003] EP-A-0,446,948 discloses a method for the immobilization of microbial cells wherein microbial cells are mixed with polyuronic acid, and a calcium chloride solution is dropwise added to the mixture so as to form polyuronate gel, thereby immobilizing the cells by the polyuronate gel, and subsequently the polyuronate gel is crosslinked with a crosslinking agent such as an allylamine polymer to prevent leakage of an enzyme.


[0004] WO 97/46590 discloses a process for immobilizing microbial cells wherein (i) a support member is coated with a hydrophilic polymeric surfactant which is then crosslinked with an excess amount of a crosslinker to form a first layer, (ii) the microbial cells are immobilized on the surface of the first layer using the excess (free) crosslinker, and (iii) the residual (free) crosslinker is used to attach a second layer, which can be an allylamine polymer, to the first layer, thereby sandwiching the cells between the first and second layers.

SUMMARY OF INVENTION


[0005] It is therefore an object of the present invention to provide a novel immobilized biocatalyst which overcomes the disadvantages of considerable inactivation of the enzyme during the immobilization.

[0006] As a result of much research aimed at overcoming the aforementioned disadvantage, the present inventors completed the present invention upon finding that an immobilized biocatalyst lacking that disadvantage and capable of stably maintaining its enzymatic activity for extended periods can be obtained by binding enzyme-containing cells to a solid support using a polyallylamine-based polymer which is water-soluble under acidic conditions but becomes insoluble under alkali conditions.

[0007] The present invention therefore provides an immobilized biocatalyst comprising enzyme-containing cells which are immobilized on a water-insoluble solid support by a polymer represented by the following general formula:



wherein Y is a direct bond or a divalent group represented by the following formula




R₁ and R₂ are each independently hydrogen atoms or organic residues, X⁻ represents an anion, and n is an integer of 100 to 5000, the immobilized biocatalyst being obtainable by mixing the said cells with water in the neutral to weakly basic liquid polymer, sprinkling the mixture onto the solid support and drying the solid support coated with the mixture.

[0008] The present invention further provides a process for producing L-aspartic acid which is characterized by contacting fumaric acid with ammonia or ammonium fumarate, or maleic acid with ammonia or ammonium maleate on an immobilized biocatalyst according to the invention which employs cells containing aspartase and/or maleate isomerase, or a combination of such cells, as the above-mentioned cells.


DETAILED DESCRIPTION

[0009] The support used for the immobilized biocatalyst of the invention may be either in granular or sheet form, with examples of granules including ion-exchange resins, for example the anionic exchange resins such as Ambalite IRA-904, IRA-94S, Ambalite A-26, A-27 (products of Organo, KK.), and the cationic exchange resins such as Ambalite 200C, 201B, IRC-50 and IRC-76 (products of Organo, KK.); and inorganic carriers such as molecular sieves, alumina, silica and silica gel. The size of the granules is preferably 0.1 mm to 10 mm, and more preferably 0.3 mm to 3 mm. As sheets there may be used ion-exchange films and alumina or silica sheets. The thickness of sheets used is preferably 0.01 mm to 10 mm, and more preferably 0.1 mm to 5 mm.

[0010] According to the invention, the resin used for immobilization of the cells is a polymer represented by the following general formula:








wherein Y is a direct bond or a divalent group represented by the following formula



R₁ and R₂ are each independently hydrogen atoms or organic residues, X⁻ represents an anion, and n is an integer of 100 to 5000.

[0011] When R₁ and R₂ in the formula are organic residues, they may be alkyl groups of 10 carbon atoms or less such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and tert-butyl, among which methyl is particularly preferred. There may also be used halogen- or hydroxyl-substituted organic residues such as 4-chloro-2-dimethylpentyl, 3-ethyl-2,5-dichloroheptyl and 2-hydroxy-3,5-dimethylnonyl, although 3-chloro-2-hydroxypropyl is particularly preferred.

[0012] X⁻ may be a halogen ion such as F⁻, Cl⁻, Br⁻ or I⁻, among which Cl⁻ is preferred. Other monovalent anions such as NO₃⁻ and the like may also be used. The number represented by n is preferably 100 to 5000. Specific polymers which may be mentioned include those represented by the following structural formulas.



30

35

40

45

50

55

[0013] An example of a polymer of formula (A) is PAS410 (tradename), an example of a polymer of formula (B) is PAS92 (tradename), an example of a polymer of formula (C) is PAS-M-1 (tradename), an example of a polymer of formula (D) is PAS-880 and an example of a polymer of formula (E) is PAS-H-5L, all of which may be obtained from Nitto Spinning, KK. Alternatively, for example, the polymer of formula (A) may be produced by 1:1 copolymerization of a diallylamine salt (e.g. a hydrochloride) and maleic acid, the polymer of formula (B) may be produced by copolymerization or radical polymerization of a diallylamine salt (e.g. a hydrochloride) and sulfur dioxide gas, and the polymer of

formula (C) may be produced by copolymerization or radical polymerization of diallylmonoalkylamines.

[0014] The cells used for the invention may be any cells containing the desired enzyme, including for example, microbial cells such as bacteria, yeast, algae and filamentous fungi, and plant cells. The enzyme is not particularly restricted, and examples of various enzymes include aspartase which converts ammonium fumarate into L-aspartic acid, maleate isomerase which converts maleic acid into fumaric acid, fumarase which converts fumaric acid into L-malic acid, malease which converts maleic acid into D-malic acid, and hydratase which converts cis-epoxy-succinic acid into L-tartaric acid. More specific examples of cells which may be used are bacterial cells containing aspartase, bacterial cells containing maleate isomerase, and bacterial cells containing both aspartase and maleate isomerase.

[0015] The following are more specific examples of microbial cells which may be used.

Table 1

Types of reactions using immobilized biocatalysts

Starting material	Product	Enzyme	Cells
Ammonium fumarate	L-aspartic acid	aspartase	<i>Escherichia coli</i> ATCC 11303 " ATCC 9637 " ATCC 27325 <i>Brevibacterium</i>
Maleic acid	fumaric acid	maleate isomerase	<i>Pseudomonas maltophilia</i> ATCC 13270
Ammonium maleate	L-aspartic acid	maleate isomerase + aspartase	<i>Alcaligenes</i> <i>Alcaligenes faecalis</i> ATCC 8750 Also, combinations of microbes with maleate isomerase activity and microbes with aspartase activity
Fumaric acid	L-malic acid	fumarase	Microbes with aspartase activity <i>Escherichia coli</i> IEO 3301 <i>Bacillus stearothermophilus</i> DSM 2234 <i>Sulfolobus solfataricus</i> ATCC 49255
Maleic acid	D-malic acid	malease	<i>Pseudomonas pseudoalcaligenes</i> NSM-5 NSM-6 FERM P-15561 P-15562
Cis-epoxy-succinic acid	L-tartaric acid	hydratase	<i>Acinetobacter tartarogenes</i> ATCC 31105

[0016] When a polymer represented by the above general formula (4) is added at low concentration to a culture solution obtained by culturing one of these microbes, the cultured cells can be condensed and precipitated, thus simplifying the procedure since centrifugation of a large amount of the culture solution is unnecessary.

[0017] Specifically, the polymer represented by the above general formula (4) is added to a culture solution of the

microbes to be immobilized to a solid content of 10-10,000 ppm, and preferably 100-1000 ppm, and after being allowed to stand the microbial cells aggregate and precipitate. Cooling the culture solution in advance is effective for preventing reduction of the desired enzyme activity.

[0018] After thus precipitating the microbial cells, the supernatant may be drawn out to reduce the liquid amount by a few times. For example, after 500 ppm of PAS-880 has been added to 1 L of culture solution and allowed to stand for 30 minutes, approximately 800 ml may be separated off as the supernatant. The remaining cell aggregate can then be subjected to low-speed centrifugation at 3000 rpm to recover the cells as a cake.

[0019] Because the aggregated cells can be sufficiently separated even with a low-speed centrifuge, it is possible to recover the cells with simple equipment.

[0020] The recovered cells may then be mixed with the pH-adjusted polymer either directly or after addition of water, and used as the immobilized material.

[0021] In immobilization of the cells, either one type of cell may be immobilized or two or more different types of cells may be immobilized. For example, bacterial cells containing maleate isomerase and bacterial cells containing aspartase may be immobilized in combination.

[0022] The cell immobilization may be accomplished by first mixing the cells with water in a neutral to weakly basic liquid polymer, and sprinkling it onto the solid support and drying it. The drying may be carried out using a common particle drier capable of not inactivating the enzyme in the cells, such as a rotary evaporator, fluidized bed drier or spray drier.

[0023] When the amount of immobilized cells is insufficient after a single operation, the sprinkling operation and drying operation of the polymer/cell mixture on the solid support may be repeated a number of times, for example 2 to 5 times. The proportion of the cells to be immobilized and the polymer is 4 to 200 kg, and preferably 10 to 50 kg, of polymer per 1 kg of cells (dry weight). When preparing a mixture of the polymer and cells, the amount of water added is 5 to 400 kg, and preferably 20 to 100 kg, per 1 kg of the cells. The thickness of the polymer layer containing the immobilized cells is 0.01 to 0.3 mm, and preferably 0.1 to 0.2 mm.

[0024] If the immobilized catalyst is in the form of a film, the above-mentioned cell/polymer mixture may be applied onto a sheet-like support and then dried. The density of the immobilized cells may be adjusted by varying the amount of application or the number of times the application/drying cycle is repeated.

[0025] One exemplified application for an immobilized biocatalyst of the invention is production of L-aspartic acid. In this case, an immobilized biocatalyst with immobilized aspartase-containing cells, especially bacterial cells, is packed into a column and an aqueous solution of fumaric acid and ammonia or an aqueous solution of ammonium fumarate is passed through the column.

[0026] Alternatively, an immobilized biocatalyst with immobilized cells (for example, bacterial cells) containing aspartase and maleate isomerase, or an immobilized catalyst with both immobilized cells containing aspartase and cells containing maleate isomerase on a single immobilizing support, is packed into a column and an aqueous solution of maleic acid and ammonia or an aqueous solution of ammonium maleate is passed through the column, to produce L-aspartic acid by a single-step process from maleic acid.

[0027] Still alternatively, both an immobilized biocatalyst with immobilized cells containing maleate isomerase and an immobilized catalyst with immobilized cells containing aspartase are combined and packed into a column and an aqueous solution of maleic acid and ammonia or an aqueous solution of ammonium maleate is passed through the column, to produce L-aspartic acid by a single-step process from maleic acid.

[0028] Still alternatively, an immobilized biocatalyst with immobilized cells containing maleate isomerase and an immobilized catalyst with immobilized cells containing aspartase are packed into separate columns, and an aqueous solution of maleic acid and ammonia or an aqueous solution of ammonium maleate is first passed through the immobilized biocatalyst with immobilized maleate isomerase to convert the maleic acid to fumaric acid, and is then passed through the column of the immobilized biocatalyst with immobilized aspartase to convert the ammonium fumarate to L-aspartic acid.

EXAMPLES

[0029] The present invention will now be explained in more detail by way of examples, although the invention is in no way to be limited thereby. The reaction products were analyzed by liquid chromatography.

Example 1

[0030] A 5 L jar fermenter was loaded with 3 L of medium (pH 6.3) comprising a composition of 10 g of fumaric acid, 5 g of $(\text{NH}_4)_2\text{SO}_4$, 1 g of KH_2PO_4 , 3 g of K_2HPO_4 , 0.5 g of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5.5 g of NaOH and 20 g of yeast extract in 1 L of distilled water, and *Escherichia coli* ATCC 11303 was inoculated thereto for aeration agitation culturing at 37°C. Culturing was continued for twenty hours, and the cells were collected by centrifugation.

[0031] After preparing a mixture of 70 g of PAS-880 (product of Nitto Spinning, KK.) with the pH adjusted to near 7 with an alkali and 230 g of deionized water, the above-mentioned cells were dispersed therein. In a 6 L volume eggplant-shaped flask was placed 300 ml of an ion-exchange resin (Ambalite IRA-94S Cl, average particle size: 0.5 mm, product of Organo, KK.) and 20 0.5-inch Teflon spheres, after which 1/6 of the previously obtained cell dispersion was added and subjected to vacuum drying for one hour with an evaporator at 30°C while rotating, to coat the cells with the ion-exchange resin. After this operation was repeated 6 times, the Teflon spheres were removed to obtain beads of immobilized biocatalyst.

[0032] The immobilized biocatalyst as prepared above was immersed overnight in a 20% ammonium fumarate solution (pH 8.5) at 4°C, after which 50 ml thereof was filled into a jacketed column and 30°C warm water was circulated through the jacket to bring the temperature of the reactor to 30°C.

[0033] A substrate solution in a capped bottle (200 g of fumaric acid, 200 g of 25% ammonia water, 0.25 g of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ and 1 g of sodium sulfite in 1 L, adjusted to pH 8.3 with ammonia) was fed through a Teflon tube and passed through the column at a rate of 25 ml per hour for continuous reaction.

[0034] Upon analyzing the reaction solution in the 6th hour after initiating the reaction, a reaction product i.e., L-aspartase acid was found in a roughly equimolar amount compared to the fumaric acid consumed, with a reaction conversion rate of 99.7%.

[0035] The conversion rate of 99.7% was maintained up to the 7th day, 1st month and even 4th month after initiating the reaction. No change was observed in the form of the immobilized biocatalyst.

Examples 2 to 4

[0036] The procedure of Example 1 was repeated using PAS-M-1 (Example 2), PAS-410 (Example 3) and PAS-H-5L (Example 4) as immobilizing polymers instead of PAS-880.

[0037] The results are summarized in Table 2.

Table 2

	Polymer	Initial conversion rate (%)	Conversion rate after 7 days (%)
Example 1	PAS-880	99.7	99.7
2	PAS-M-1	99.8	99.7
3	PAS-410	99.8	99.7
4	PAS-H-5L	99.6	99.6

[0038] The conversion rate from fumaric acid to L-aspartic acid did not decrease even after 7 days of use.

Example 5

[0039] *Escherichia coli* ATCC 11303 was cultured, in the same manner as Example 1, by aeration/agitation culturing at 37°C. After 20 hours of culturing the medium was cooled to 10°C, 3 g of PAS-880 was added, and after 5 minutes of agitation the mixture was left to stand without agitation. When the cells aggregated and precipitated after 30 minutes, about 2.5 L of the supernatant was drawn out. The remaining solution was then drawn out together with the cells and subjected to low-speed centrifugation to recover 120 g of wet cells.

[0040] After preparing a mixture of 70 g of PAS-880 (product of Nitto Spinning, KK.) with the pH adjusted to near 7 with an alkali and 170 g of deionized water, the above-mentioned cells were dispersed therein. In a 6 L volume eggplant-shaped flask were placed 300 ml of an ion-exchange resin (Ambalite IRA-94S Cl, average particle size: 0.5 mm, product of Organo, KK.) and 20 0.5-inch Teflon spheres, after which 1/6 of the previously obtained cell dispersion was added and subjected to vacuum drying for one hour with an evaporator at 30°C while rotating, to coat the cells with the ion-exchange resin. After this operation was repeated 6 times, the Teflon spheres were removed to obtain beads of an immobilized biocatalyst.

[0041] The immobilized biocatalyst as prepared above was immersed overnight in a 20% ammonium fumarate solution (pH 8.5) at 4°C, after which 50 ml thereof was filled into a jacketed column and 30°C warm water was circulated through the jacket to bring the temperature of the reactor to 30°C.

[0042] A substrate solution in a capped bottle (200 g of fumaric acid, 200 g of 25% ammonia water, 0.25 g of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ and 1 g of sodium sulfite in 1 L, adjusted to pH 8.3 with ammonia) was fed through a Teflon tube and passed through the column at a rate of 25 ml per hour for continuous reaction.


[0043] Upon analyzing the reaction solution in the 6th hour after initiating the reaction, a reaction product i.e., L-

aspartase acid was found in a roughly equimolar amount with the fumaric acid consumed, with a reaction conversion rate of 99.7%.


[0044] The conversion rate of 99.7% was maintained up to the 7th day, 1st month and even 4th month after initiating the reaction. No change was observed in the form of the immobilized biocatalyst.

Claims

1. An immobilized biocatalyst comprising enzyme-containing cells which are immobilized on a water-insoluble solid support by a polymer represented by the following general formula:



wherein Y is a direct bond or a divalent group represented by the following formula:




R_1 and R_2 are each independently hydrogen atoms or organic residues, X^- represents an anion, and n is an integer of 100-5000, the immobilized biocatalyst being obtainable by mixing the said cells with water in the neutral to weakly basic liquid polymer, sprinkling the mixture onto the solid support and drying the solid support coated with the mixture.


2. An immobilized biocatalyst according to claim 1, wherein said support is an ion-exchange resin or inorganic support.
3. An immobilized biocatalyst according to claim 1 or 2, wherein said support comprises granules or a sheet.
4. An immobilized biocatalyst according to any of claims 1 to 3, wherein R_1 and R_2 in the formula of the polymer are each independently hydrogen or alkyl of up to 10 carbon atoms, optionally substituted by hydroxyl or halogen, and X^- is a halogen or NO_3^- ion.
5. An immobilized biocatalyst according to any of claims 1 to 4, which comprises cells containing enzymes selected from fumarase, malease, hydratase, aspartase and maleate isomerase.
6. An immobilized biocatalyst according to any of claims 1 to 4, which comprises bacterial cells containing aspartase and/or maleate isomerase, or a combination of such bacterial cells.
7. The use of an immobilized biocatalyst according to any of claims 1 to 6 for the manufacture of fumaric acid, L-aspartic acid, L-malic acid, D-malic acid or L-tartaric acid.
8. A process for producing L-aspartic acid or fumaric acid, **characterized by** reacting fumaric acid with ammonia or ammonium fumarate, or maleic acid with ammonia or ammonium maleate, or maleic acid on an immobilized biocatalyst according to claim 6.

Patentansprüche


1. Immobilisierter Biokatalysator aufweisend Enzym enthaltend Zellen, die an einem wasserunlöslichen festen Träger immobilisiert sind mittels eines Polymers, das dargestellt wird durch die folgende allgemeine Formel:



worin Y eine direkte Bindung oder eine zweiwertige Gruppe ist, die dargestellt wird durch die folgende Formel:



oder




R₁ und R₂ jeweils unabhängig voneinander Wasserstoffatome oder organische Reste sind, X[⊖] für ein Anion steht und n eine ganze Zahl von 100 bis 5000 ist, wobei der immobilisierte Biokatalysator erhältlich ist durch Mischen der Zellen mit Wasser in das neutrale bis schwach basische flüssige Polymer, Spritzen des Gemisches auf den festen Träger und Trocknen des mit dem Gemisch beschichteten festen Trägers.


2. Immobilisierter Biokatalysator nach Anspruch 1, bei dem der Träger ein Ionenaustauscherharz oder ein anorganischer Träger ist.
3. Immobilisierter Biokatalysator nach Anspruch 1 oder 2, bei dem der Träger Granulat oder ein Flachmaterial aufweist.
4. Immobilisierter Biokatalysator nach einem der Ansprüche 1 bis 3, bei dem R₁ und R₂ in der Formel des Polymers jeweils unabhängig voneinander Wasserstoff oder Alkyl mit bis zu 10 Kohlenstoffatomen, gewünschtenfalls mit Hydroxyl oder Halogen substituiert, sind und X[⊖] ein Halogen- oder NO₃[⊖]-Ion ist.
5. Immobilisierter Biokatalysator nach einem der Ansprüche 1 bis 4, der Zellen aufweist, die Enzyme enthalten, welche ausgewählt sind unter Fumarase, Malease, Hydratase, Aspartase und Maleatisomerase.
6. Immobilisierter Biokatalysator nach einem der Ansprüche 1 bis 4, der Bakterienzellen, die Aspartase und/oder Maleatisomerase enthalten, oder eine Kombination derartiger Bakterienzellen aufweist.
7. Verwendung eines immobilisierten Biokatalysators nach einem der Ansprüche 1 bis 6 zur Herstellung von Fumarsäure, L-Asparaginsäure, L-Äpfelsäure, D-Äpfelsäure oder L-Weinsäure.
8. Verfahren zur Herstellung von L-Asparaginsäure oder Fumarsäure, **gekennzeichnet durch** zur Reaktion Bringen von Fumarsäure mit Ammoniak oder Ammoniumfumarat, oder von Maleinsäure mit Ammoniak oder Ammoniummaleat, oder von Maleinsäure an einem immobilisierten Biokatalysator nach Anspruch 6.

Revendications

1. Biocatalyseur immobilisé comprenant des cellules contenant des enzymes qui sont immobilisées sur un support solide insoluble dans l'eau par un polymère répondant à la formule générale suivante :



dans laquelle Y est une liaison directe ou un groupe divalent répondant à la formule suivante :



R₁ et R₂ sont chacun indépendamment des atomes d'hydrogène ou des résidus organiques, X⁻ représente un anion et n est un nombre entier de 100 à 5000, le biocatalyseur immobilisé pouvant être obtenu en mélangeant ces cellules avec de l'eau dans le polymère liquide neutre à faiblement basique, en pulvérisant le mélange sur le support solide et en séchant le support solide revêtu du mélange.

2. Biocatalyseur immobilisé selon la revendication 1, dans lequel ce support est une résine échangeuse d'ions ou un support minéral.
3. Biocatalyseur immobilisé selon la revendication 1 ou la revendication 2, dans lequel ce support comprend des granules ou une feuille.
4. Biocatalyseur immobilisé selon l'une quelconque des revendications 1 à 3, dans lequel le R₁ et le R₂ de la formule du polymère sont chacun indépendamment l'hydrogène ou un groupe alkyle ayant jusqu'à 10 atomes de carbone, substitué si on le désire par un groupe hydroxyle ou un halogène et X⁻ est un halogène ou un ion NO₃⁻.
5. Biocatalyseur immobilisé selon l'une quelconque des revendications 1 à 4, qui comprend des cellules contenant des enzymes choisies parmi la fumarase, la maléase, l'hydratase, l'aspartase et la maléate isomérase.
6. Biocatalyseur immobilisé selon l'une quelconque des revendications 1 à 4, qui comprend des cellules bactériennes contenant de l'aspartase et/ou de la maléate isomérase, ou une combinaison de ces cellules bactériennes.
7. Utilisation d'un biocatalyseur immobilisé selon l'une quelconque des revendications 1 à 6 pour la préparation de l'acide fumarique, de l'acide L-aspartique, de l'acide L-malique, de l'acide D-malique ou de l'acide L-tartrique.
8. Procédé de préparation de l'acide L-aspartique ou de l'acide fumarique, caractérisé en ce que l'on fait réagir l'acide fumarique avec de l'ammoniaque ou le fumarate d'ammonium, ou l'acide maléique avec l'ammoniaque ou le maléate d'ammonium, ou l'acide maléique sur un biocatalyseur immobilisé selon la revendication 6.